

Bounded and unitary elements in pro- C^* -algebras ^{*}

Rachid El Harti and Gábor Lukács [†]

February 2, 2008

Abstract

A *pro- C^* -algebra* is a (projective) limit of C^* -algebras in the category of topological $*$ -algebras. From the perspective of non-commutative geometry, pro- C^* -algebras can be seen as non-commutative k -spaces. An element of a pro- C^* -algebra is *bounded* if there is a uniform bound for the norm of its images under any continuous $*$ -homomorphism into a C^* -algebra. The $*$ -subalgebra consisting of the bounded elements turns out to be a C^* -algebra. In this paper, we investigate pro- C^* -algebras from a categorical point of view. We study the functor $(-)_b$ that assigns to a pro- C^* -algebra the C^* -algebra of its bounded elements, which is the dual of the Stone-Ćech-compactification. We show that $(-)_b$ is a coreflector, and it preserves exact sequences. A generalization of the Gelfand duality for commutative unital pro- C^* -algebras is also presented.

1. Introduction

In a C^* -algebra, the norm is determined by the algebraic structure, and thus being a C^* -algebra is an entirely algebraic property. This intuitive conclusion was confirmed by van Osdol, who proved that the unit ball functor, sending a C^* -algebra A to the set $\{a \in A \mid \|a\| \leq 1\}$, is monadic (cf. [35]). Later on, Pelletier and Rosický investigated the system of operations and equations representing the unit balls of C^* -algebras as universal algebras (cf. [26] and [25]).

A Hausdorff k -space is a colimit of compact Hausdorff spaces in the category of Hausdorff spaces (and their continuous maps). Such spaces were thoroughly investigated by Brown, Steenrod, and Dubuc and Porta (cf. [9, 3.3], [34], and [12]). A *pro- C^* -algebra* is a (projective) limit of C^* -algebras in the category of topological $*$ -algebras. Such algebras were studied under various names (LMC^* -algebras, locally C^* -algebras, and σ - C^* -algebra in the metrizable case) by Schmüdgen [32], Inoue [16], Arveson [4], Phillips [27], and El Harti [14].

By the Gelfand duality, the category of commutative unital C^* -algebras is equivalent to the opposite of the category of compact Hausdorff spaces. It turns out that using a close relative of k -spaces, the Gelfand duality can be extended to commutative unital pro- C^* -algebras (Theorem 3.7).

^{*}2000 Mathematics Subject Classification: 18A05 46H05 46J05 46K05

[†]The second author gratefully acknowledges the generous financial support received from the Killam Trusts, and Dalhousie University that enabled him to do this research.

Thus, from the perspective of non-commutative geometry, pro- C^* -algebras can be seen as non-commutative k -spaces.

An element a of a pro- C^* -algebra A is *bounded* if there is a constant M such that $\|\varphi(a)\| \leq M$ for every continuous $*$ -homomorphism $\varphi: A \rightarrow B$ into a C^* -algebra B . The smallest constant M with this property is denoted by $\|a\|_\infty$, and the $*$ -subalgebra A_b of bounded elements in A is a C^* -algebra with respect to $\|\cdot\|_\infty$ (see section 4). Since the Stone-Ćech-compactification of a Tychonoff space X can be obtained as the space of multiplicative functionals of the C^* -algebra $C_b(X)$ with respect to the norm of uniform convergence, it is natural to think of the functor $A \mapsto A_b$ as the non-commutative dual of the Stone-Ćech-compactification functor $X \mapsto \beta X$.

In this paper, we investigate pro- C^* -algebras from a categorical point of view. After a brief presentation of terminology and basic results on pro- C^* -algebras in section 2, we proceed by presenting a generalization of the Gelfand duality for commutative unital pro- C^* -algebras in section 3. We also compare this generalized Gelfand duality with results of Dubuc and Porta [12]. In section 4, we study the functor $(-)_b$ that assigns to a pro- C^* -algebra A the C^* -algebra A_b of its bounded elements. We show that this functor is a coreflector (Theorem 4.8), and it preserves exactness of sequences (Theorem 4.11). Finally, in section 5, we make a few observations concerning the connected component of the identity of the group of unitary elements in a pro- C^* -algebra.

2. Preliminaries

In this section, we introduce some terminology, and present well-known elementary results on pro- C^* -algebras.

2.1. Topological $*$ -algebras. A $*$ -algebra (or *involutive algebra*) is an algebra A over $\mathbb{C}$ with an involution $*$: $A \rightarrow A$ such that $(a + \lambda b)^* = a^* + \bar{\lambda}b^*$ and $(ab)^* = b^*a^*$ for every $a, b \in A$ and $\lambda \in \mathbb{C}$. A linear seminorm p on a $*$ -algebra A is a C^* -seminorm if $p(ab) \leq p(a)p(b)$ (*submultiplicative*) and $p(a^*a) = p(a)^2$ (C^* -condition) for every $a, b \in A$. Note that the C^* -condition alone implies that p is submultiplicative, and in particular $p(a^*) = p(a)$ for every $a \in A$ (cf. [33], [24, 9.5.14]). A *topological $*$ -algebra* is a $*$ -algebra A equipped with a topology making the operations (addition, multiplication, additive inverse, involution) jointly continuous. The category of topological $*$ -algebras and their continuous $*$ -homomorphisms is denoted by T^*A . For $A \in T^*A$, one puts $\mathcal{N}(A)$ for the set of continuous C^* -seminorms on A ; $\mathcal{N}(A)$ is a directed set with respect to pointwise ordering, because $\max\{p, q\} \in \mathcal{N}(A)$ for every $p, q \in \mathcal{N}(A)$.

2.2. C^* -algebras. A C^* -algebra is a complete Hausdorff topological algebra whose topology is given by a single C^* -norm. The full subcategory of T^*A formed by the C^* -algebras is denoted by C^*A . For $A \in T^*A$ and $p \in \mathcal{N}(A)$, $\ker p = \{a \in A \mid p(a) = 0\}$ is a $*$ -ideal in A , and p induces a C^* -norm on the quotient $A/\ker p$, so the completion A_p of this quotient with respect to p is a C^* -algebra. Each pair $p, q \in \mathcal{N}$ such that $q \geq p$ induces a natural (continuous) surjective $*$ -homomorphism $\pi_{pq}: A_q \rightarrow A_p$, which turns $A_{(-)}: \mathcal{N}(A) \rightarrow C^*A$ into a functor.

2.3. The $*$ -representation topology. For $A \in \mathsf{T}^*A$, a $*$ -representation of A is a continuous $*$ -homomorphism $\pi: A \rightarrow B(H)$ of A into the C^* -algebra of bounded operators on some Hilbert space H (i.e., a morphism in T^*A). The class of $*$ -representations of A is denoted by $\mathcal{R}(A)$. By the Gelfand-Naïmark-Segal theorem, every C^* -algebra is $*$ -isomorphic (and thus isometric) to a closed subalgebra of $B(H)$ for a large enough Hilbert space H (cf. [11, 2.6.1]), so each A_p embeds (isometrically) into $B(H_p)$ for some Hilbert space H_p . Thus, for each $p \in \mathcal{N}(A)$ one obtains a $*$ -representation $\bar{\pi}_p: A \rightarrow A/\ker p \rightarrow A_p \rightarrow B(H_p)$ such that $p(x) = \|\bar{\pi}_p(x)\|$. Conversely, each $\pi \in \mathcal{R}(A)$ gives rise to a C^* -seminorm $p_\pi(x) = \|\pi(x)\|$. Therefore, the initial topology $\mathcal{T}_A$ induced by the class $\mathcal{R}(A)$ coincides with the one induced by the family of C^* -seminorms $\mathcal{N}(A)$. The topology $\mathcal{T}_A$ is called the $*$ -representation topology.

2.4. Pro- C^* -algebras. A pro- C^* -algebra is a complete Hausdorff topological $*$ -algebra A whose topology is defined by a family of C^* -seminorms, or equivalently, the topology of A coincides with $\mathcal{T}_A$, and $\mathcal{T}_A$ is Hausdorff and complete (i.e., every Cauchy filter or net converges). For $A \in \mathsf{T}^*A$, the following statements are equivalent (cf. [28, 1.1.1]):

- (i) $A \cong \varprojlim_{p \in \mathcal{N}(A)} A_p$ in T^*A ;
- (ii) A is a (projective) limit of C^* -algebras in T^*A ;
- (iii) A is a closed $*$ -subalgebra of a product of C^* -algebras in T^*A ;
- (iv) A is a pro- C^* -algebra.

It turns out that for a pro- C^* -algebra A , $(A/\ker p, p)$ is already complete, so A_p is a quotient of A (cf. [2, Thm. 2.4], [32, Folg. 5.4], [28, 1.2.8], [27, 1.12]). Pro- C^* -algebras form a full reflective subcategory of T^*A that we denote by $\overline{\mathsf{P}^*A}$ (cf. [20, 5.1]). In light of (i), we often refer to an element a of a pro- C^* -algebra A as $(a_p)_{p \in \mathcal{N}(A)}$, where $a_p \in A_p$ and $\pi_{pq}(a_q) = a_p$ for every $p, q \in \mathcal{N}(A)$.

2.5. Examples.

- (A) $\prod_{n=1}^{\infty} \mathbb{M}_n(\mathbb{C})$, the product of the (full) matrix algebras $\mathbb{M}_n(\mathbb{C})$ in T^*A , is a metrizable pro- C^* -algebra.
- (B) The algebra $C_\omega([0, 1])$ of continuous complex-valued maps on $[0, 1]$ with the topology of uniform convergence on compact countable subsets is a pro- C^* -algebra (for details, see section 3).
- (C) The tangent algebra of a C^* introduced by Arveson (defined as the universal factorizer of derivations into C^* -algebras) fails to be C^* -algebra, but it is a metrizable pro- C^* -algebra (for details, see [4, 5.9] and [28, 2.1]).

2.6. Spectrum and unitization. We denote by G_A the set of invertible elements in an algebra A . The spectrum of an element a in a unital algebra A over $\mathbb{C}$ is defined as

$$\mathrm{sp}_A(a) = \{\lambda \in \mathbb{C} \mid a - \lambda 1 \notin G_A\}. \quad (1)$$

If $A \in \overline{\mathsf{P}^*A}$, then $a = (a_p) \in A$ is invertible if and only if each a_p is invertible in A_p . Therefore,

$$\mathrm{sp}_A(a) = \bigcup_{p \in \mathcal{N}(A)} \mathrm{sp}_{A_p}(a_p). \quad (2)$$

If $B \subseteq A$ is a closed $*$ -subalgebra that contains the unit of A and $b \in B$, then $\text{sp}_B(b) = \text{sp}_A(b)$ (cf. [32, 1.2]). If A is not unital (which is the case for ideals in a pro- C^* -algebra), one can adjoin a unit by putting $A^+ = A \oplus \mathbb{C}e$, extending multiplication in the obvious way, and setting $(a, \lambda)^* = (a^*, \bar{\lambda})$. (Certainly, $A^+ = \varprojlim A_p^+$ is a pro- C^* -algebra.) In this case, one defines $\text{sp}_{A^+}(a)$ as the spectrum of a . This definition is consistent (up to $\{0\}$), because if A already had a unit, then $\text{sp}_{A^+}(a) = \text{sp}_A(a) \cup \{0\}$. Thus, if I is a closed (two-sided) $*$ -ideal of a unital pro- C^* -algebra A and $c \in I$, then $\text{sp}_{I^+}(c) = \text{sp}_{A^+}(c) = \text{sp}_A(c)$, because c cannot be invertible in A .

2.7. Functional calculus. Let A be a unital pro- C^* -algebra and let $a = (a_p) \in A$. Suppose that f is a continuous complex-valued map on $\text{sp}_A(a)$ and either a is *normal* (i.e., $a^*a = aa^*$) or f is analytic on a neighborhood of $\text{sp}_A(a)$. It follows from (2) that f is continuous on each $\text{sp}_{A_p}(a_p)$, and either each a_p is normal or f is analytic on a neighborhood of each $\text{sp}_{A_p}(a_p)$. In both cases, $f(a_p)$ is defined, $\text{sp}_{A_p}(f(a_p)) = f(\text{sp}_{A_p}(a_p))$, and $\pi_{pq}(f(a_q)) = f(a_p)$ for each pair $q \geq p$ in $\mathcal{N}(A)$. Therefore, $(f(a_p)) \in A$ and $\text{sp}_A(f(a)) = f(\text{sp}_A(a))$. In case A is not unital, $f(0) = 0$ is the only additional assumption needed in order to ensure that $f(a)$ (calculated in A^+) belongs to A .

2.8. Automatic continuity fails in $\overline{\mathbf{P}^*\mathbf{A}}$. It is a well-known fact that every $*$ -homomorphism of an involutive Banach algebra into a C^* -algebra is continuous (furthermore, its norm is ≤ 1 ; cf. [11, 1.3.7]). In particular, every $*$ -homomorphism between C^* -algebras is continuous. Results of this nature are referred to as “automatic continuity” of homomorphisms, because the continuity is deduced from a purely algebraic assumption (cf. [23, Chapter 6]). Unfortunately, in general, this result fails for pro- C^* -algebras: Both $C(\omega_1)$ and $C(\omega_1 + 1)$ are pro- C^* -algebras in the topology of uniform convergence on compacta, but the $*$ -homomorphism $C(\omega_1) \rightarrow C(\omega_1 + 1)$ given by setting $f(\omega_1) = \lim_{x \rightarrow \omega_1} f(x)$ fails to be continuous (cf. [22, 12.2], [28, 1.4.9]).

2.9. Automatic continuity for σ - C^* -algebras. A σ - C^* -algebra is a pro- C^* -algebra whose topology is metrizable, or equivalently, it is a (projective) limit of a sequence of C^* -algebras in $\mathbf{T}^*\mathbf{A}$. Every $*$ -homomorphism of a σ - C^* -algebra into a pro- C^* -algebra is continuous (cf. [27, 5.2], [28, 1.1.6]).

3. Commutative pro- C^* -algebras

By the Gelfand duality, every commutative unital C^* -algebra is $*$ -isomorphic (and thus isometrically isomorphic) to the algebra of continuous maps on the set of its multiplicative functionals (i.e., characters). In this section, we present some generalizations of this result to pro- C^* -algebras. Our aim is to adhere to a categorical point of view, and thus it slightly differs from the approach of Inoue [16] and Phillips [28, 1.4] & [27, 2], who obtained essentially the same results.

3.1. Leading example. For a Hausdorff space X , one puts $\mathcal{K}(X)$ for the collection of compact subsets of X . For $\mathcal{F} \subseteq \mathcal{K}(X)$ such that $\bigcup \mathcal{F} = X$, we say that a map $f: X \rightarrow Y$ is $\mathcal{F}$ -continuous if $f|_F$ is continuous for every $F \in \mathcal{F}$. We put $C_{\mathcal{F}}(X)$ for the $*$ -algebra of complex-valued $\mathcal{F}$ -continuous maps f on X , and provide it with the topology of *uniform convergence on $\mathcal{F}$* given

by the family of C^* -seminorms $\{p_F\}_{F \in \mathcal{F}}$, where $p_F(f) = \sup_{x \in F} |f(x)|$. It is a well-known result on uniform spaces that $C_{\mathcal{F}}(X)$ is complete (cf. [17, 7.10(d)]), and thus $C_{\mathcal{F}}(X)$ is a pro- C^* -algebra. We say that $\mathcal{F}$ is a *strongly functionally generating* family for X if every map in $C_{\mathcal{F}}(X)$ is continuous (cf. [3, p. 111]). The Gelfand duality states that every commutative unital C^* -algebra is isomorphic in C^*A (and thus in T^*A) to the algebra $C(K)$ of continuous maps with the topology of uniform convergence on some compact Hausdorff space K . We show that every commutative unital pro- C^* -algebra is isomorphic to $C_{\mathcal{F}}(X)$ in T^*A for some space X , where X is strongly functionally generated by $\mathcal{F}$.

3.2. The categories $k\text{Haus}$ and $k_R\text{Tych}$. A Hausdorff space X is a k -space if every $\mathcal{K}(X)$ -continuous (i.e., k -continuous) map $f: X \rightarrow Y$ into a Hausdorff space Y is continuous, or equivalently, if $U \subseteq X$ is open provided that $U \cap K$ is open in K for every compact subset K of X (cf. [15, 3.3.21]). The category $k\text{Haus}$ of Hausdorff k -spaces and continuous maps is cartesian closed, and it is a coreflective subcategory of the category Haus of Hausdorff spaces (cf. [9, 3.3], [21, VII.8]). (The coreflector $k: \text{Haus} \rightarrow k\text{Haus}$ is often referred to as k -ification. Some authors use the term “compactly generated spaces” for k -spaces.) A Hausdorff space X is a k_R -space if it is strongly functionally generated by $\mathcal{K}(X)$ (i.e., every real-valued k -continuous map $f: X \rightarrow \mathbb{R}$ on X is continuous). Lukács showed that the category $k_R\text{Tych}$ of Tychonoff k_R -spaces and continuous maps is cartesian closed, $k_R\text{Tych}$ is a coreflective subcategory of Tych (Tychonoff spaces), and $k_R\text{Tych}$ is equivalent to a full epireflective subcategory of $k\text{Haus}$ (cf. [19, Theorems 3.2, 2.1, 3.1]). The motivation for introducing this category is the incompatibility of the k -space property and the Tychonoff property (cf. [19, Example]). Their necessity for pro- C^* -algebras roots in the observation that if X admits a strongly functionally generating family of compact subsets, then it must be a k_R -space.

3.3. The character space $\Delta(A)$. For a topological vector space X , one puts X' for the *dual space* of X consisting of the continuous linear functionals on X . The w^* -topology on X' is the topology of pointwise convergence on the points of X —in other words, X' is provided with the initial topology with respect to all evaluations maps $\{\text{ev}_x: X' \rightarrow \mathbb{C}\}_{x \in X}$. Since (X', w^*) is a Hausdorff topological vector space, it is Tychonoff. For $A \in T^*A$, one puts $\Delta(A)$ for the subspace of (A', w^*) consisting of the non-zero $*$ -homomorphisms. Using this notation, Gelfand duality states that the map $\text{ev}_A: A \rightarrow C(\Delta(A))$, defined by $a \mapsto \text{ev}_a$, is an isomorphism of C^* -algebras for every C^* -algebra A .

3.4. Theorem. *Let A be a commutative unital pro- C^* -algebra. Then:*

- (a) $\Delta(A) = \bigcup_{p \in \mathcal{N}(A)} \Delta(A_p)$;
- (b) $\Delta(A)$ is a Tychonoff k_R -space, and $\Phi(A) = \{\Delta(A_p) \mid p \in \mathcal{N}(A)\}$ is a strongly functionally generating family;
- (c) the natural map $\text{ev}_A: A \rightarrow C_{\Phi(A)}(\Delta(A))$ defined by $a \mapsto \text{ev}_a$ is an isomorphism of pro- $*$ -algebras.

Proof. (a) By 3.3, $\Delta(A)$ is a subspace of the Tychonoff space A' , and so it is Tychonoff. Since A is commutative and unital, so are its quotients A_p . Thus, $A_p \cong C(\Delta(A_p))$ by Gelfand duality,

and the $\Delta(A_p)$ are compact. Each project $A \rightarrow A_p$ gives rise to a continuous injective map $\Delta(A_p) \rightarrow \Delta(A)$, which is an embedding, because $\Delta(A_p)$ is compact. On the other hand, if $\chi \in \Delta(A)$, then $p_\chi(a) = |\chi(a)|$ is a continuous C^* -seminorm on A , and therefore $\chi \in \Delta(A_{p_\chi})$.

(b) Suppose that $f: \Delta(A) \rightarrow \mathbb{R}$ is $\Phi(A)$ -continuous, in other words, $f|_{\Delta(A_p)}$ is continuous for every $p \in \mathcal{N}(A)$. Then, by Gelfand duality, for each $p \in \mathcal{N}(A)$ there is $a_p \in A_p$ such that $f|_{\Delta(A_p)} = \text{ev}_{a_p}$. Furthermore, since $f|_{\Delta(A_q)}|_{\Delta(A_p)} = f|_{\Delta(A_p)}$, one has $\pi_{pq}(a_q) = a_p$ for every $q \geq p$. Thus, $a = (a_p) \in A$ and $f = \text{ev}_a$. In particular, f is continuous, and therefore $\Phi(A)$ is a strongly functionally generating family. Hence, $\Delta(A)$ is a k_R -space.

(c) It is clear that ev_A is a $*$ -homomorphism. Since

$$p(a) = \sup\{|\rho(a)| \mid \rho \in \Delta(A_p)\} = p_{\Delta(A_p)}(\text{ev}_a) \quad (3)$$

for every $p \in \mathcal{N}(A)$, it follows that ev_A is continuous and open onto its image. If $\text{ev}_A(a) = 0$ for $a \in A$, then $\rho(a) = 0$ for every $\rho \in \Delta(A)$, and so $p(a) = 0$ for every $p \in \mathcal{N}(A)$. Thus, $a = 0$ (because A is Hausdorff). If $f \in C_{\Phi(A)}(\Delta(A))$, then by the argument developed in (b), $f = \text{ev}_a$ for some $a \in A$. Therefore, ev_A is surjective. Hence, ev_A is an isomorphism of pro- C^* -algebras. $\square$

3.5. Lemma. *Let X be a Tychonoff k_R -space, and let $\mathcal{F}$ be a strongly functionally generating family of compact subsets of X . Then the natural map $\text{ev}_X: X \rightarrow \Delta(C_{\mathcal{F}}(X))$ defined by $x \mapsto \text{ev}_x$ is a homeomorphism.*

Proof. Since X is a Tychonoff space k_R -space, it embeds into $C_{\mathcal{F}}(X)'$ by evaluation (cf. [19, Lemma 2.1]), and its image is contained in $\Delta(C_{\mathcal{F}}(X))$, so ev_X is an embedding. In order to show that ev_X is surjective, let $\rho \in \Delta(C_{\mathcal{F}}(X))$. Since ρ is a continuous functional with respect to the topology determined by the seminorms $\{p_F\}_{F \in \mathcal{F}}$, there are $F_1, \dots, F_l \in \mathcal{F}$ and a constant r such that $|\rho(f)| \leq r \max\{p_{F_1}(f), \dots, p_{F_l}(f)\}$ for every $f \in C_{\mathcal{F}}(X)$. Thus, for $F_0 = F_1 \cup \dots \cup F_l$, ρ factors through a multiplicative functional $\bar{\rho}$ of the C^* -algebra $C(F_0)$, that is, $\rho(f) = \bar{\rho}(f|_{F_0})$. Therefore, by Gelfand duality, there is $x_0 \in F_0$ such that $\bar{\rho}(g) = g(x_0)$ for every $g \in C(F_0)$, and hence $\rho = \text{ev}_X(x_0)$, as desired. $\square$

3.6. Notations. The category SFG is defined as follows: Objects are pairs $(X, \mathcal{F})$, where X is a Tychonoff k_R -space and $\mathcal{F}$ is a strongly functionally generating family of compact subsets of X . A morphism $g: (X, \mathcal{F}) \rightarrow (Y, \mathcal{G})$ in SFG is a continuous map with the property that for every $F \in \mathcal{F}$, there is a finite subfamily $\{G_1, \dots, G_l\} \subseteq \mathcal{G}$ such that $g(F) \subseteq G_1 \cup \dots \cup G_l$. We denote by $C_1 \overline{P^*A}$ the full subcategory of $\overline{P^*A}$ formed by the commutative unital pro- C^* -algebras.

3.7. Theorem. *The pair of functors*

$$C_1 \overline{P^*A}^{\text{op}} \longrightarrow \text{SFG} \quad (4)$$

$$A \longmapsto (\Delta(A), \Phi(A)) \quad (5)$$

$$C_{\mathcal{F}}(X) \longleftarrow (X, \mathcal{F}) \quad (6)$$

form an equivalence of categories.

Proof. By Theorem 3.4(c), $A \cong C_{\Phi(A)}(\Delta(A))$ for every $A \in C_1\overline{P^*A}$. On the other hand, by Lemma 3.5, $X \cong \Delta(C_{\mathcal{F}}(X))$ for every $(X, \mathcal{F}) \in \text{SFG}$. Furthermore, $F \in \Phi(C_{\mathcal{F}}(X))$ for every $F \in \mathcal{F}$, because $p_F \in \mathcal{N}(C_{\mathcal{F}}(X))$. In order to complete the proof, let $G \in \Phi(C_{\mathcal{F}}(X))$ (and we show that it is contained in a finite union of members of $\mathcal{F}$). Then $p_G \in \mathcal{N}(C_{\mathcal{F}}(X))$, and thus there are $F_1, \dots, F_l \in \mathcal{F}$ and a constant r such that $p_G \leq r \max\{p_{F_1}, \dots, p_{F_l}\}$, because $\{p_F\}_{F \in \mathcal{F}}$ generates the topology of $C_{\mathcal{F}}(X)$. Since X is Tychonoff, if $x_0 \notin F_1 \cup \dots \cup F_l$, then there exists a continuous map h on X such that $h(x_0) = 1$ and $h|_{F_i} = 0$ for each i . So, $p_{F_i}(h) = 0$, and therefore $p_G(h) = 0$. In particular, $x_0 \notin G$, and hence $G \subseteq F_1 \cup \dots \cup F_l$, as desired. $\square$

3.8. k_R -ification using function spaces. Recall that for a Tychonoff space X , the Stone-Čech compactification βX (i.e., the reflection of X into the category of compact Hausdorff spaces) can be computed using Gelfand duality: $\beta X \cong \Delta(C_b(X))$, where $C_b(X)$ is the C^* -algebra of all continuous bounded maps on X with the norm of uniform convergence. Theorem 3.7 provides a similar method for finding the k_R -ification $k_R X$ of a Tychonoff space X (that is, the coreflection of X into $k_R \text{Tych}$): By [19, Lemma 2.1(b)], $k_R X \cong \Delta(C_{\mathcal{K}(X)}(X))$. Naturally, when X is not a k_R -space, $C_{\mathcal{K}(X)}(X)$ contains some non-continuous functions that are only k -continuous on X .

3.9. K -algebras. Dubuc and Porta investigated the category of $*$ -algebra objects in $k\text{Haus}$, called K -algebras (cf. [12]). These are $*$ -algebras on a Hausdorff k -space A , with continuous unitary operations, and with binary operations being continuous on $A \times_k A = k(A \times A)$ (which is the cartesian product in $k\text{Haus}$). So, addition and multiplication are expected to be only k -continuous on $A \times A$. A typical example of a K -algebra is kA , the k -ification of a pro- C^* -algebra A . Dubuc and Porta established a dual adjunction between commutative unital K -algebras and *functionally Hausdorff* k -spaces (i.e., continuous real-valued maps separate points) using the cotensor of the $k\text{Haus}$ -enriched category of K -algebras, and showed that the Gelfand duality is a restriction of this dual adjunction. They also proved that the k -ification of a commutative pro- C^* -algebra admits a canonical monad-algebra structure with respect to the resulting monad (cf. [12, 3.8]). An essential difference between this setting and the one present in this paper is that in the context of K -algebras, the natural topology on $\Delta(A)$ is the k -ified compact-open topology, and not the w^* -topology.

Our approach is akin to the methods employed by Dubuc and Porta in a later paper, and by Phillips, using *quasitopological spaces* (cf. [13] and [27]). For instance, Phillips showed that $C_1\overline{P^*A}^{\text{op}}$ is equivalent to the category QHaus_f of functionally Hausdorff quasitopological spaces (cf. [27, 2.7]). Thus, by Theorem 3.7, we conclude:

3.10. Corollary. *The categories SFG and QHaus_f are equivalent.* $\square$

4. The algebra of bounded elements

For a Tychonoff k_R -space X , the C^* -algebra $C_b(X)$ of continuous complex-valued bounded maps on X is precisely the subalgebra of $C(X)$ of the maps $f: X \rightarrow \mathbb{C}$ such that $\sup_{K \in \mathcal{K}(X)} p_K(f) < \infty$. In

this section, a natural generalization of this notion to the non-commutative case is presented.

4.1. Bounded elements. For a pro- C^* -algebra A , $\|a\|_\infty = \sup_{p \in \mathcal{N}(A)} p(a)$ is called the *uniform norm*, and $a \in A$ is *bounded* if $\|a\|_\infty < \infty$. The $*$ -subalgebra A_b of the bounded elements in A is dense in A , and A_b is a C^* -algebra with the norm $\|\cdot\|_\infty$ (cf. [32, Satz 3.1], [28, 1.2.4, 1.2.7], [27, 1.11]). One way to view A_b is to observe that it is $\varprojlim_{p \in \mathcal{N}(A)} A_p$ in C^*A . While A carries the subspace topology induced by the product topology $\prod_{p \in \mathcal{N}(A)} A_p$, A_b is equipped with the topology of “uniform convergence,” as expected. Thus, the inclusion $(A_b, \|\cdot\|_\infty) \rightarrow A$ is continuous. If $B \subseteq A$ is a closed $*$ -algebra of A , then $B_b = A_b \cap B$ (cf. [32, 3.4]).

4.2. Spectrally bounded elements. The *spectral radius* of an element a in a pro- C^* -algebra A is $r_A(a) = \sup\{|\lambda| \mid \lambda \in \text{sp}_A(a)\} = \sup_{p \in \mathcal{N}(A)} r_{A_p}(a_p)$, and a is *spectrally bounded* if $r_A(a) < \infty$. (If A is not unital, sp_{A^+} is used instead of sp_A .) We put A_{sb} for the set of spectrally bounded elements in A (it fails to be a subalgebra). For every $a \in A$, $r_A(a) \leq \|a\|_\infty$ holds, and so $A_b \subseteq A_{sb}$. However, if a is normal, then $p(a) = r_{A_p}(a)$, and so $\|a\|_\infty = r_A(a)$. In particular, a normal element is bounded if and only if it is spectrally bounded. Therefore, A_b is spanned by the spectrally bounded self-adjoint elements in A (i.e., for $A_s = \{a \in A \mid a = a^*\}$, $A_{sb} \cap A_s$ spans A_b), because each $a \in A$ can be written as $a = a_1 + ia_2$ with $a_1, a_2 \in A_s$. Since the spectrum of an element does not depend on the topology of A , we obtain:

4.3. Corollary. *For a pro- C^* -algebra A , A_b depends only on the algebraic structure of the underlying $*$ -algebra of A . In other words, if A_1 and A_2 are pro- C^* -algebras with the same underlying $*$ -algebra (i.e., only the topologies differ), then $(A_1)_b = (A_2)_b$. $\square$*

4.4. Example for $A_{sb} \neq A_b$. In the σ - C^* -algebra $A = \prod_{n=1}^{\infty} \mathbb{M}_n(\mathbb{C})$ (see 2.5(A)), consider the quasi-nilpotent element $L = (L_n)$ given by

$$L_{n+1} = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 2 & 0 & \cdot & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & n-1 & 0 \\ 0 & 0 & \dots & 0 & 0 & n \\ 0 & 0 & \dots & 0 & 0 & 0 \end{pmatrix}. \quad (7)$$

Since each L_n is nilpotent, $\text{sp}_{M_n(\mathbb{C})}(L_n) = \{0\}$, and so L is spectrally bounded. But, on the other hand,

$$n \leq \sqrt{r_{M_{n+1}(\mathbb{C})}(L_{n+1}L_{n+1}^*)} \leq \|L\|_\infty, \quad (8)$$

which implies that L cannot be bounded. Therefore, $A_{sb} \neq A_b$.

4.5. Example for $A_{sb} = A$. Let $X = [0, 1]$, and put $\mathcal{D}$ for the collection of compact countable subsets of X . Since X is metrizable, $\mathcal{D}$ is a strongly functionally generating family, and thus $A = C_{\mathcal{D}}([0, 1])$ is a pro- C^* -algebra that is not a C^* -algebra. Clearly, $A_{sb} = A_b = A$ (as sets), nevertheless, A is not a C^* -algebra. This also shows that $C_{\mathcal{D}}([0, 1])$ is not a σ - C^* -algebra:

4.6. Proposition. *If A is a σ - C^* -algebra such that every element of A is spectrally bounded, then A is a C^* -algebra.*

Proof. The inclusion $(A_b, \|\cdot\|_\infty) \rightarrow A$ is always continuous. Since $A_{sb} = A$, one has $A_b = A$, because A_b is spanned by $A_{sb} \cap A_s$. Thus, the identity map $A \rightarrow (A_b, \|\cdot\|_\infty)$ is a $*$ -homomorphism from a σ - C^* -algebra into a C^* -algebra, and therefore continuous by 2.9. $\square$

4.7. The functor $(-)_b$. If $\varphi: A \rightarrow B$ is a $*$ -homomorphism of $*$ -algebras, then $\varphi(A_s) \subseteq B_s$ and $\varphi(A_{sb}) \subseteq B_{sb}$, because $\text{sp}_B(\varphi(a)) \subseteq \text{sp}_A(a)$ for every $a \in A$. Thus, $\varphi(A_s \cap A_{sb}) \subseteq B_s \cap B_{sb}$. Therefore, if A and B are pro- C^* -algebras, then $\varphi(A_b) \subseteq B_b$. Hence, φ induces a $*$ -homomorphism $\varphi_b: A_b \rightarrow B_b$. Since no assumptions were made about the continuity of φ , this shows that $(-)_b: \overline{\mathbf{P}^* \mathbf{A}_d} \rightarrow \mathbf{C}^* \mathbf{A}$ is a functor, where $\overline{\mathbf{P}^* \mathbf{A}_d}$ is the category of pro- C^* -algebras and their (not necessarily continuous) $*$ -homomorphisms (cf. [32, Folg. 3.3] and [27, 1.13]).

4.8. Theorem. $(-)_b: \overline{\mathbf{P}^* \mathbf{A}_d} \rightarrow \mathbf{C}^* \mathbf{A}$ is a coreflector.

Proof. Let B be a C^* -algebra, A be a pro- C^* -algebra, and $\varphi: B \rightarrow A$ be a $*$ -homomorphism. Then $\varphi_b: B_b \rightarrow A_b$ is a $*$ -homomorphism, and therefore φ factors through A_b uniquely, because $B_b = B$. $\square$

4.9. Corollary. *Let A be a unital pro- C^* -algebra and $a \in A$ be normal. Then $\|f(a)\|_\infty \leq \|f\|_\infty$ for every $f \in C(\text{sp}_A(a))$. In particular, if f is bounded, then so is $f(a)$.* $\square$

4.10. Exact sequences in $\overline{\mathbf{P}^* \mathbf{A}_d}$. A sequence

$$\cdots \longrightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} C \longrightarrow \cdots \quad (9)$$

in $\overline{\mathbf{P}^* \mathbf{A}_d}$ is *exact at B* if $\ker \beta = \alpha(A)$. A categorical explanation for this choice of definition of exactness is that it is inherited from $\mathbf{T}^* \mathbf{A}_d$, the category of topological $*$ -algebras with (not necessarily continuous) $*$ -homomorphisms. See also Remark 4.12.

4.11. Theorem. *The functor $(-)_b: \overline{\mathbf{P}^* \mathbf{A}_d} \rightarrow \mathbf{C}^* \mathbf{A}$ preserves exactness of sequences.*

Proof. Consider the sequence in (9). Since $\beta\alpha = 0$, one has $\beta_b\alpha_b = 0$, and so $\alpha_b(A_b) \subseteq \ker \beta_b$. Observe that $\alpha_b(A_b)$ is closed in B_b , because α_b is a $*$ -homomorphism between C^* -algebras. Thus, in order to show the reverse inclusion, it suffices to show that every self-adjoint $b \in \ker \beta_b$ is the limit of a sequence from $\alpha_b(A_b)$. Every $b \in \ker \beta_b$ has the form $b = \alpha(a)$ for some $a \in A$ because (9) is exact at B . If b is self-adjoint, then by replacing a with $\frac{a + a^*}{2}$ if necessary, we may assume that a is self-adjoint too. By Corollary 4.9, $f_n(a) \in A_b$ for

$$f_n(x) = \frac{n^2 x}{n^2 + x^2}. \quad (10)$$

Since $f_n(x) \rightarrow x$ uniformly on $[-\|b\|_\infty, \|b\|_\infty]$, one has $\alpha_b(f_n(a)) = f_n(b) \xrightarrow{\|\cdot\|_\infty} b$. (Observe that $f_n(a)$ is a rational polynomial in a , which means that $\alpha(f_n(a))$ is determined by $\alpha(a)$ even if α is not continuous). Therefore, $b \in \alpha_b(A_b)$, as desired. $\square$

4.12. Remark. The notion of exactness provided in 4.10 might appear counterintuitive at first sight, because the $*$ -homomorphic image $\alpha(A)$ need not be complete (e.g., $B = C_{\mathcal{K}(\mathbb{R})}(\mathbb{R})$, $A = C_b(\mathbb{R})$, and $\alpha = \text{id}$). Requiring $\overline{\alpha(A)} = \ker \beta$ instead could make the impression of being a better definition, however, this is not the case: For $A = C([0, 1])$, $B = \mathbb{C}^{[0,1]}$, and α the inclusion, $\overline{\alpha(A)}$ is dense in B , but $\alpha_b(A)$ is a closed proper subalgebra of $B_b = l^\infty([0, 1])$. Therefore, $\alpha(A) = \ker \beta$ is indeed necessary in Theorem 4.11.

4.13. Corollary. *Let I be a closed $*$ -ideal of the pro- C^* -algebra A such that A/I is complete (i.e., A/I is a pro- C^* -algebra). Then I_b is a closed $*$ -ideal of A_b , and $(A/I)_b \cong A_b/I_b$.*

Proof. Apply Theorem 4.11 to the sequence $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$. □

4.14. The quotient A/I need not be complete. Because of the automatic continuity property of σ - C^* -algebras (see 2.9), every short exact sequence of σ - C^* -algebras is the limit of a countable collection of short exact sequences of C^* -algebras (cf. [28, 5.3]). In particular, if A is a σ - C^* -algebra and I is its closed $*$ -ideal, then A/I is complete (cf. [28, 5.4]). The quotient of a complete topological algebra by a closed ideal need not be complete (cf. [18]), but one would hope that pro- C^* -algebras are better behaved. This is, however, not the case, as the following example shows: Let X be a locally compact space that is not normal. Then X contains a closed subspace $Y \subseteq X$ such that not every continuous map on Y can be extended continuously to X . Put $A = C_{\mathcal{K}(X)}(X)$ and $B = C_{\mathcal{K}(Y)}(Y)$, let $\varphi: A \rightarrow B$ be the restriction (i.e., $f \mapsto f|_Y$), and set $I = \ker \varphi$. Since Y is closed in X , it is easy to see that φ is open onto its image, and thus $A/I \cong \varphi(A)$ as topological $*$ -algebras. The image, $\varphi(A)$, is precisely the $*$ -subalgebra of continuous maps on Y that can be continuously extended to X , and so $\varphi(A) \neq B$. On the other hand, by the Stone-Weierstrass Theorem, $\varphi(A)$ is dense in B . Therefore, $\varphi(A) \cong A/I$ is not complete. (This example, which is based on [31, Exercise IV.11], was suggested by Subhash Bhatt, and was communicated personally to the authors by N. C. Phillips.)

4.15. Corollary. *Let A be a pro- C^* -algebra. Then $A_p \cong A_b/(\ker p)_b$ for every $p \in \mathcal{N}(A)$.* □

4.16. Proposition. *Let A be a pro- C^* -algebra. If A_b is simple, then A is a C^* -algebra.*

Proof. Since A_b is dense in A (see 4.1), $(\ker p)_b \neq A_b$ for every nonzero $p \in \mathcal{N}(A)$. Thus, $(\ker p)_b = 0$, because A_b is simple, and therefore $\ker p = 0$, because $(\ker p)_b$ is dense in $\ker p$. Hence, by Corollary 4.15, $A = A_p = A_b$ for every nonzero $p \in \mathcal{N}(A)$, which means that A is a C^* -algebra. □

4.17. Problem. Let B be a unital C^* -algebra that is not simple. Is there a pro- C^* -algebra A that is not a C^* -algebra such that $B = A_b$? This problem is fairly non-trivial even in the abelian case, where $B = C(K)$ for some compact Hausdorff space K . If K is metrizable and uncountable, then $A = C_{\mathcal{D}}(K)$ is a positive solution, where $\mathcal{D}$ is the collection of compact countable subsets of K . In fact, it is sufficient to assume that K is compact and a *sequential space* (i.e., continuity of maps is determined by continuity on sequences). Nevertheless, we do not have an answer for the problem even for such well-known C^* -algebras as c and c_0 (the subalgebras of l^∞ consisting of the convergent and zero sequences, respectively).

5. The group of unitary elements

In this section, we are concerned with the topological group A_u of unitary elements in a unital pro- C^* -algebra A (i.e., $xx^* = x^*x = 1$), and study the connected component $A_{u,1}$ of 1 in it. A key tool in our investigation is the subgroup $A_{\text{exp}} = \{e^{ia_1} \cdots e^{ia_n} \mid a_1, \dots, a_n \in A_s\}$, generated by exponentials of self-adjoint elements.

5.1. Proposition. *Let A be a pro- C^* -algebra. Then:*

- (a) A_{exp} is path-connected in the topology of A ;
- (b) $A_{\text{exp}} \subseteq A_{u,1}$;
- (c) A_{exp} is an clopen subgroup of A_u in the $\|\cdot\|_\infty$ -topology.

Proof. We note that since a unitary element has norm 1 in any C^* -algebra, $\|u\|_\infty = 1$ for every $u \in A_u$. So $A_u \subseteq A_b$, and $A_u = (A_b)_u$ as sets (their topology might be quite different, though).

(a) We show that each generator e^{ia} ($a \in A_s$) can be joined to 1 by a path. To that end, consider the map $u: [0, 1] \rightarrow A$ given by $u(t) = e^{ita}$. It suffices to show that u is continuous. For each $p \in \mathcal{N}(A)$, $f(t, s) = e^{its}$ is uniformly continuous on the compact set $[0, 1] \times \text{sp}_{A_p}(a_p)$, and thus $u_p(t) = e^{ita_p}$ is continuous $[0, 1] \rightarrow A_p$. Therefore, u is continuous, as desired.

(b) follows from (a).

(c) Since every open subgroup in a topological group is also closed, it suffices to show that $\{u \in A_u \mid \|1-u\|_\infty < 1\} \subseteq A_{\text{exp}}$. Let $u \in A_u$ be such that $\|1-u\|_\infty < 1$. Then $\text{sp}_A(u) \subseteq \mathbb{S} \setminus \{-1\}$ (where $\mathbb{S} = \{z \in \mathbb{C} \mid |z| = 1\}$), and so $f(z) = \arg z$ can be continuously defined on $\text{sp}_A(u)$. Thus, by the functional calculus, $u = e^{ia}$ for $a = f(u)$, and $a \in A_s$ because f is real-valued. Therefore, $u \in A_{\text{exp}}$, as desired. $\square$

5.2. Remark. If A is a C^* -algebra, then A_{exp} is a clopen path-connected subgroup of A_u (by Proposition 5.1(c)), and hence $A_{\text{exp}} = A_{u,1}$.

5.3. Lemma. *Let $\alpha: A \rightarrow B$ be continuous $*$ -homomorphism of pro- C^* -algebras. Then one has $\alpha(g(a)) = g(\alpha(a))$ for every normal $a \in A$ and $g \in C(\text{sp}_A(a))$.*

Proof. Put $b = \alpha(a)$. For every polynomial $p(z, \bar{z})$, $\alpha(p(a, a^*)) = p(b, b^*)$, because α is a $*$ -homomorphism. If $g \in C(\text{sp}_A(a))$, then it can be uniformly approximated by polynomials $p_\alpha(z, \bar{z})$ on compact sets, and therefore $g(a) = \lim p_\alpha(a, a^*)$ and $g(b) = \lim p_\alpha(b, b^*)$. Hence, the statement follows by continuity of α . $\square$

5.4. Corollary. *Let $\alpha: A \rightarrow B$ be a surjective continuous $*$ -homomorphism of pro- C^* -algebras. Then $\alpha(A_{\text{exp}}) = B_{\text{exp}}$.* $\square$

5.5. Theorem. *Let A be a pro- C^* -algebra. Then:*

- (a) A_{exp} is dense in $A_{u,1}$;
- (b) $A_{u,1} = \varprojlim_{p \in \mathcal{N}(A)} (A_p)_{u,1}$.

Proof. To shorten notations, put $G = \varprojlim_{p \in \mathcal{N}(A)} (A_p)_{u,1}$. Clearly, $A_{u,1} \subseteq G \subseteq A_u$.

(a) We show that $A_{\exp}$ is dense in G . Since each projection $\pi_p: A \rightarrow A_p$ is surjective, by Corollary 5.4, $\pi_p(A_{\exp}) = (A_p)_{\exp}$. Since A_p is a C^* -algebra, $(A_p)_{\exp} = (A_p)_{u,1}$ (cf. Remark 5.2), and thus $\pi_p(A_{\exp}) = (A_p)_{u,1}$. Therefore, the statement follows, because $\mathcal{N}(A)$ is directed.

(b) Since $A_{\exp} \subseteq A_{u,1}$, $A_{u,1}$ is dense in G . On the other hand, $A_{u,1}$ is a closed subgroup of A_u , because it is a connected component. Therefore, $A_{u,1} = G$, as desired. $\square$

5.6. Example for $A_{\exp} \neq A_{u,1}$. Let $B = C([0, 1]^\omega) \otimes M_2(\mathbb{C})$ be the C^* -algebra of continuous maps $f: [0, 1]^\omega \rightarrow M_2(\mathbb{C})$ from the Hilbert cube into $M_2(\mathbb{C})$ (or equivalently, of 2×2 matrices with entries in $C([0, 1]^\omega)$). Phillips showed that for every $n \in \mathbb{N}$, there is $u_n \in B_u$ in the *path* component of 1 in B_u such that u_n cannot be expressed as the product of n many exponentials e^{ib} , where $b \in B_s$ (cf. [30, 2.6]). In other words, $u_n \neq e^{ib_1} \cdots e^{ib_n}$ for every $b_1, \dots, b_n \in B_s$. Put $A = B^\omega$, the countable power of B in T^*A ; A is a σ - C^* -algebra, and $u = (u_n) \in A_u$. If $p_n: \mathbb{I} \rightarrow B_u$ is a path such that $p_n(0) = 1$ and $p_n(1) = u_n$, then $p = (p_n): \mathbb{I} \rightarrow A_u$ is a path joining u to 1 in A_u . Thus, u is in the path component of A_u , but it cannot be in $A_{\exp}$, because u is not a product of n exponentials for any $n \in \mathbb{N}$. In particular, $A_{\exp} \subsetneq A_{u,1}$. (This example is due to N. C. Phillips [30, 2.11].)

Acknowledgments

We wish to thank Professor N. C. Phillips for the valuable discussions that were of great assistance in writing this paper.

References

- [1] Graham R. Allan. Stable inverse-limit sequences, with application to Fréchet algebras. *Studia Math.*, 121(3):277–308, 1996.
- [2] Constantin Apostol. b^* -algebras and their representation. *J. London Math. Soc. (2)*, 3:30–38, 1971.
- [3] A. V. Arkhangel'skiĭ. *Topological function spaces*, volume 78 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1992. Translated from the Russian by R. A. M. Hoksbergen.
- [4] William Arveson. The harmonic analysis of automorphism groups. In *Operator algebras and applications, Part I (Kingston, Ont., 1980)*, volume 38 of *Proc. Sympos. Pure Math.*, pages 199–269. Amer. Math. Soc., Providence, R.I., 1982.
- [5] Subhash J. Bhatt and Dinesh J. Karia. An intrinsic characterization of Pro- C^* -algebras and its applications. *J. Math. Anal. Appl.*, 175(1):68–80, 1993.
- [6] Michael E. Boardman. Relative spectra in complete lmc-algebras with applications. *Illinois J. Math.*, 39(1):119–139, 1995.
- [7] G. Bourdaud. Quelques aspects de la dualité en analyse fonctionnelle. *Diagrammes*, 5:B1–B12, 1981.
- [8] Gérard Bourdaud. Sur la dualité des algèbres localement convexes. *C. R. Acad. Sci. Paris Sér. A-B*, 281(23):Ai, A1011–A1014, 1975.
- [9] R. Brown. Function spaces and product topologies. *Quart. J. Math. Oxford Ser. (2)*, 15:238–250, 1964.

- [10] M. Chidami and R. El Harti. Calcul fonctionnel holomorphe en dimension infinie dans les lmc. *Rend. Circ. Mat. Palermo* (2), 48(3):541–548, 1999.
- [11] Jacques Dixmier. C^* -algebras. North-Holland Publishing Co., Amsterdam, 1977. Translated from the French by Francis Jellet, North-Holland Mathematical Library, Vol. 15.
- [12] Eduardo J. Dubuc and Horacio Porta. Convenient categories of topological algebras, and their duality theory. *J. Pure Appl. Algebra*, 1(3):281–316, 1971.
- [13] Eduardo J. Dubuc and Horacio Porta. Uniform spaces, Spanier quasitopologies, and a duality for locally convex algebras. *J. Austral. Math. Soc. Ser. A*, 29(1):99–128, 1980.
- [14] Rachid El Harti. Contractible Fréchet algebras. *Proc. Amer. Math. Soc.*, 132(5):1251–1255 (electronic), 2004.
- [15] Ryszard Engelking. *General topology*, volume 6 of *Sigma Series in Pure Mathematics*. Heldermann Verlag, Berlin, second edition, 1989. Translated from the Polish by the author.
- [16] Atsushi Inoue. Locally C^* -algebra. *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 25:197–235, 1971.
- [17] John L. Kelley. *General topology*. D. Van Nostrand Company, Inc., Toronto-New York-London, 1955.
- [18] Gottfried Köthe. *Topological vector spaces. I*. Translated from the German by D. J. H. Garling. Die Grundlehren der mathematischen Wissenschaften, Band 159. Springer-Verlag New York Inc., New York, 1969.
- [19] Gábor Lukács. A convenient subcategory of Tych. *Appl. Categ. Structures*, 12(4):369–377, 2004.
- [20] Gábor Lukács. Lifted closure operators. *Preprint*, ArXiv: math.CT/0502410.
- [21] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [22] Ernest A. Michael. Locally multiplicatively-convex topological algebras. *Mem. Amer. Math. Soc.*, 1952(11):79, 1952.
- [23] Theodore W. Palmer. *Banach algebras and the general theory of $*$ -algebras. Vol. I*, volume 49 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1994. Algebras and Banach algebras.
- [24] Theodore W. Palmer. *Banach algebras and the general theory of $*$ -algebras. Vol. 2*, volume 79 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2001. $*$ -algebras.
- [25] J. Wick Pelletier and J. Rosický. On the equational theory of C^* -algebras. *Algebra Universalis*, 30(2):275–284, 1993.
- [26] Joan Wick Pelletier and Jiří Rosický. Generating the equational theory of C^* -algebras and related categories. In *Categorical topology and its relation to analysis, algebra and combinatorics (Prague, 1988)*, pages 163–180. World Sci. Publishing, Teaneck, NJ, 1989.
- [27] N. Christopher Phillips. Inverse limits of C^* -algebras. *J. Operator Theory*, 19(1):159–195, 1988.
- [28] N. Christopher Phillips. Inverse limits of C^* -algebras and applications. In *Operator algebras and applications, Vol. 1*, volume 135 of *London Math. Soc. Lecture Note Ser.*, pages 127–185. Cambridge Univ. Press, Cambridge, 1988.
- [29] N. Christopher Phillips. Representable K -theory for σ - C^* -algebras. *K-Theory*, 3(5):441–478, 1989.
- [30] N. Christopher Phillips. How many exponentials? *Amer. J. Math.*, 116(6):1513–1543, 1994.
- [31] H. H. Schaefer and M. P. Wolff. *Topological vector spaces*, volume 3 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1999.
- [32] Konrad Schmüdgen. Über LMC-Algebren. *Math. Nachr.*, 68:167–182, 1975.
- [33] Z. Sebestyén. Every C^* -seminorm is automatically submultiplicative. *Period. Math. Hungar.*, 10(1):1–8, 1979.

- [34] N. E. Steenrod. A convenient category of topological spaces. *Michigan Math. J.*, 14:133–152, 1967.
- [35] D. H. Van Osdol. C^* -algebras and cohomology. In *Categorical topology (Toledo, Ohio, 1983)*, volume 5 of *Sigma Ser. Pure Math.*, pages 582–587. Heldermann, Berlin, 1984.
- [36] Wiesław Żelazko. *Banach algebras*. Elsevier Publishing Co., Amsterdam-London-New York, 1973. Translated from the Polish by Marcin E. Kuczma.

Department of Mathematics
University Hassan I, FST de Settat
BP 577, 2600 Settat
Morocco

e-mail: elharti@ibnsina.uh1.ac.ma

Department of Mathematics and Statistics
Dalhousie University
Halifax, B3H 3J5, Nova Scotia
Canada

e-mail: lukacs@mathstat.dal.ca